UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

International General Certificate of Secondary Education

MARK SCHEME for the May/June 2012 question paper for the guidance of teachers

0580 MATHEMATICS

0580/23

Paper 2 (Extended), maximum raw mark 70

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2012 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

| Page 2 | Mark Scheme: Teachers' version | Syllabus | Paper |
|--------|--------------------------------|----------|-------|
| | IGCSE – May/June 2012 | 0580 | 23 |

Abbreviations

cao correct answer only cso correct solution only

dep dependent

ft follow through after error isw ignore subsequent working

oe or equivalent SC Special Case

www without wrong working

soi seen or implied

| Qu | Answers | Mark | Part marks |
|-------|---------------------------|------|---|
| 1 | 95 | 2 | B1 for 85 seen or M1 $x = 180$ – their angle <i>ADC</i> , if it is clearly seen |
| 2 | 120 | 2 | M1 for $\frac{750 \times 2 \times 8}{100}$ oe seen or SC1 870 as final answer |
| 3 (a) | 3.26077 | 1 | seen |
| (b) | 3.261 | 1ft | their (a) to 4 significant figures |
| 4 | <i>y</i> ∅ −1.25 | 2 | M1 inequality with y's and constants correctly collected |
| 5 | 33 cao www | 2 | M1 any two of 5.5, 9.5, 12.5 seen |
| 6 | 31.7 | 2 | $\mathbf{M1}\ 0.5 \times 9 \times 15 \times \sin 28$ |
| 7 | u = 24(.0), v = 0.6 | 2 | B1 each |
| 8 | 7 cao | 3 | B1 for 39.5(0) or 31.5(0) or 42 M1 for (their 39.5 – 8) ÷ 4.5 or (their 42 – 10.5) ÷ 4.5 |
| 9 | $\frac{a(2-t)}{3}$ cao oe | 3 | M1 correct re-arrangement to isolate the term in w M1 correct multiplication by a M1 correct division by their 3 An incorrect answer scores a maximum of M2 |
| 10 | 10 | 3 | M1 T = $k\sqrt{l}$ A1 for $k=2$ |
| 11 | 17.05 cao www | 4 | M1 for $280 \times (1 + \frac{3}{100})^2$ oe |
| | | | M1 subtracting 280 from $280(1 + \frac{k}{100})^2$ any k |
| | | | A1 for 17.052 or SC2 297.05 on answer line |

| Page 3 | Mark Scheme: Teachers' version | Syllabus | Paper |
|--------|--------------------------------|----------|-------|
| | IGCSE – May/June 2012 | 0580 | 23 |

| 12 | (a) | $\frac{11}{12} - \frac{4}{12} \text{ oe} \\ \frac{7}{12} \text{ cao ww 0}$ | 2 | M1 correct use of a common denominator A1 |
|----|------------|--|------|--|
| | (b) | $\frac{1}{4} \times \frac{13}{11} \text{oe}$ $\frac{13}{44} \text{cao ww 0}$ | 2 | M1 inversion and operation change A1 |
| 13 | (a) | 71 | 2 | M1 for 7×8 – 3×–5 or B1 56 and –15 |
| | (b) | 3v (u + 3w) final answer | 2 | B1 for $3(uv + 3vw)$ or $v(3u + 9w)$ As final answer |
| 14 | (a) | $64p^3q^6$ | 2 | B1 $64p^{\mathrm{u}}q^{\mathrm{v}}$ or $kp^{3}q^{6}$ |
| | (b) | $0.5x^{-2} \text{ or } \frac{1}{2x^2} \text{ oe}$ | 2 | B1 $\frac{1}{2x^u}$ oe or $\frac{1}{kx^2}$ oe |
| 15 | | -3.44, 0.44 | 4 | B1 for $\sqrt{(6)^2 - 4(2)(-3)}$ or better seen |
| | | correct working must be shown | | B1 if in form $\frac{p+(or-)q}{r}$, for $p=-6$ and $r=2\times 2$ oe |
| | | | | B1 , B1 (SC1 –3.4 or –3.436 and 0.4 or 0.436) |
| 16 | | 359 www | 4 | M1 $\pi \times 4^2$ or $\frac{1}{2}\pi \times 4^2$ M1 for $0.5 \times \pi \times 8 \times 15$ oe M1 for 8×15 + their 2 ends + their curved surface area |
| 17 | (a) | (4 10) | 2 | B1 each element or correct without brackets |
| | (b) | $\begin{bmatrix} \frac{1}{2} \begin{pmatrix} 3 & -4 \\ -1 & 2 \end{bmatrix} \text{ oe }$ | 2 | B1 for $\frac{1}{2} \begin{pmatrix} a & c \\ b & d \end{pmatrix}$ or $k \begin{pmatrix} 3 & -4 \\ -1 & 2 \end{pmatrix}$ seen |
| 18 | (a) | $\mathbf{p} - \frac{1}{3} \mathbf{q}$ oe | 2 | M1 $\overrightarrow{QR} + \overrightarrow{RX}$ oe or $-\mathbf{q} + \mathbf{p} + (\frac{2}{3})\mathbf{q}$ oe |
| | (b) | $\frac{1}{2}\mathbf{p} + \frac{5}{6}\mathbf{q}$ oe | 2 ft | ft $\mathbf{q} + \frac{1}{2}$ their (a) but must be vectors or $\mathbf{M1}$ for $\overrightarrow{OQ} + \overrightarrow{QM}$ oe |
| 19 | | 6(.00) www | 4 | M1 use of area = distance M1 complete, correct set of area statements, ignoring units M1 changing min to hours or km/h to km/min |

| Page 4 | Mark Scheme: Teachers' version | Syllabus | Paper |
|--------|--------------------------------|----------|-------|
| | IGCSE – May/June 2012 | 0580 | 23 |

| 20 | $\frac{x+4}{x(x-5)}$ oe cao | 5 | B2 $(x-5)(x+4)$ seen or SC1 $(x+a)(x+b)$ where $ab = -20$ or $a+b=-1$ |
|--------|-----------------------------|---|--|
| | | | B2 $x(x-5)(x-5)$ or B1 one of $x(x^2-10x+25)$, $(x-5)(x-5)$, $(x-5)(x^2-5x)$ seen |
| 21 (a) | 7.55 www | 3 | M2 $(\frac{1}{2}\sqrt{(8^2+8^2)})^2 + 5^2$ or $4^2 + 5^2 + 4^2$ seen or M1 $8^2 + 8^2$ or $5^2 + 4^2$ or $4^2 + 4^2$ or $5^2 + (\text{their } MB)^2$ seen |
| (b) | 41.5 www | 3 | $\mathbf{M2} \sin(B) = \frac{5}{(a)} \text{ or } \tan(B) = \frac{5}{\text{their } MB} \text{ or}$ $\cos(B) = \frac{\text{their } MB}{(a)}$ or $\mathbf{M1}$ recognition of angle PBM |